

Artikel Bela

by gugus mutu mipa

Submission date: 08-Nov-2023 01:50AM (UTC-0600)

Submission ID: 2221160140

File name: Artikel_Bela.docx (856.83K)

Word count: 3245

Character count: 18711

SUPER (a, d) - $P_2 \odot P_m$ -ANTIMAGIC TOTAL LABELING OF CORONA PRODUCT OF TWO PATHS

Bela Zainun Yatin^{1*}, Zata Yumni Awanis¹, I Gede Adhitya Wisnu Wardhana¹

¹ Department of Mathematics, Faculty of Mathematics and Natural Sciences, Universitas Mataram,
 Jalan Majapahit No. 62, Mataram 83125, Indonesia

ABSTRACT

Article History:

Received : D-M-20XX
 Revised : D-M-20XX
 Accepted : D-M-20XX
 Online : D-M-20XX

Keyword:

(a, d) -H-antimagic total labeling;
 Corona product;
 H-covering;
 Path;
 Super (a, d) -H-antimagic total labeling.



This study investigates the super (a, d) - $P_2 \odot P_m$ -antimagic total labeling of the corona product $P_n \odot P_m$, where n and m are both greater than or equal to 3. We define the labeling functions for vertices and edges based on the partitioning of labels into three subsets. Using k -balanced and (k, δ) -anti balanced multisets, we demonstrate that for m being odd, $P_n \odot P_m$ is super $(9m^2n + 4mn + m - n + 3, 1)$ - $P_2 \odot P_m$ -antimagic, and for m being even, $P_n \odot P_m$ is super $(9m^2n + 4mn + m - 2n + 5, 3)$ - $P_2 \odot P_m$ -antimagic. The labeling schemes are illustrated through examples, providing insights into the antimagic properties of corona products.



<https://doi.org/10.31764/jtam.vXiY.ZZZ>



This is an open access article under the CC-BY-SA license

A. INTRODUCTION

The field of graph theory is a branch of mathematics that centers on the examination of abstract structures referred to as graphs. These graphs have been widely applied across multiple domains, including computer science, social science, and transportation (Bača et al., 2019; Chang et al., 2019; Liu et al., 2020). Their origins can be traced back to Leonard Euler's work in 1736. Within graph theory, a particularly fascinating subject of study revolves around graph labeling, which involves creating mappings from sets of vertices, edges, or both to a set of natural number (Diestel, 2017; Indunil & Perera.A.A.I., 2022; Nurvazly et al., 2022). At present, the investigation of magical and antimagical labelings continues to be a significant and actively researched area (Gallian, 2022).

The notion of magic labeling was first introduced by Sedlacek in 1963 (J. Sedlacek, 1963). This concept was subsequently expanded upon by Kotzig and Rosa in 1970 (Kotzig & Rosa, 1970) and Enomoto et al. in 1998, who introduced edge-magic total labeling and super edge-magic total labeling (Enomoto et al., 1998). Additionally, Gutiérrez and Lladó in 2005 further extended these ideas to the terms of H-(super) magic total labeling. (Gutiérrez & Lladó, 2005).

This paper focuses on a finite and simple graph G where vertex set and edge set are denoted by $V(G)$ and $E(G)$, respectively. A set of distinct subgraphs $H_1, H_2, \dots, H_k$ of G is defined as an *edge-covering* of G if every edge of G is included in at least one of the H_i . When each subgraph is isomorphic to a specific graph H , the graph G is said to have an *H-covering*. Now let G has an *H-covering*. If there exists a bijective function $f: V(G) \cup E(G) \rightarrow \{1, 2, \dots, |V(G)| + |E(G)|\}$ and a positive integer k_f such that for all subgraphs H' that are isomorphic to H , the *H-weight* $\omega(H') = \sum_{v \in V(H')} f(v) + \sum_{e \in E(H')} f(e) = k_f$, then G is called an *H-magic*. The function f is known as an *H-magic total labeling* of G . Furthermore, the labeling f is known as an *H-supermagic total labeling* of G , if the vertices of G are labeled with $1, 2, \dots, |V(G)|$. In this case, graph G is *H-supermagic* (Gutiérrez & Lladó, 2005). When H is an edge, we say that G is *(super) edge-magic* (Enomoto et al., 1998; Kotzig & Rosa, 1970). In their paper, Gutiérrez and Lladó presented star-supermagic and path-supermagic total labelings for certain connected graphs. They also offered constructions for infinite families of *H-magic* graphs for any given graph H . Furthermore, (Llado A & Moragas J, 2007) and (Ngurah et al., 2010) investigated cycle-supermagic total labellings for certain connected graphs. The readers are referred (Anjaneyulu et al., 2015; Hendy, Sugeng, & Salman, 2018; Hendy, Sugeng, Salman, et al., 2018; Sandariria et al., 2017; Simanihuruk et al., 2021; Lakshmi S & Sagayakavitha D, 2018; Ulfatimah et al., 2017; Agustin, Susanto, et al., 2019; Murugan & Chandra Kumar, 2019) for other results about *H-(super) magic* total labelling.

On the other hand, the concept of (a, d) -edge-antimagic total labeling was initially introduced by (Simanjuntak et al., 2000). This concept was later extended into the concept of *(super) (a, d)-H-antimagic total labeling* by (Inayah et al., 2009). Let a, d , and t be three positive integers, where t denotes the number of subgraphs of G that isomorphic to H . Suppose G has an *H-covering*. Graph G is called an (a, d) -*H-antimagic* if there exists a bijective function $f: V(G) \cup E(G) \rightarrow \{1, 2, \dots, |V(G)| + |E(G)|\}$ so that for all subgraphs H' that are isomorphic to H , the *H-weight* $\omega(H') = \sum_{v \in V(H')} f(v) + \sum_{e \in E(H')} f(e)$ forms an arithmetic sequence $a, a + d, a + 2d, \dots, a + (t - 1)d$. In this case, the labeling f is called an (a, d) -*H-antimagic total labeling* of G . If the vertices of G are labeled with $1, 2, \dots, |V(G)|$, then f is called the *super (a, d)-H-antimagic total labeling* of G and G is called the *super (a, d)-H-antimagic*. The readers are referred to (Martin Bača et al., 2006; Agustin et al., 2019; Hussain et al., 2012.; Inayah et al., 2013; Palanivelu & Neela, 2019; Permata Sari et al., 2019; Prihandini & Adawiyah, 2023; Raheem et al., 2014; Taimur et al., 2018) for other results about super (a, d) -*H-antimagic*.

In this paper, we study the super (a, d) -*H-antimagic* total labeling of corona product of two paths. Recall that *corona product* of two graphs G and G' , denoted by $G \odot G'$, is a graph obtained by taking one copy of G and $|V(G)|$ copies G' and then connecting the i -th vertex of G to every vertex in the i -th copy G' (Hasni et al., 2022; Permata Sari et al., 2019; Sandariria et al., 2017). We show that the corona product $P_n \odot P_m$, $m, n \geq 3$, is super (a, d) - $P_2 \odot P_m$ -antimagic where $d = 1$ if m is odd and $d = 3$ if m is even.

B. METHODS

The method used in this research is deductive method. This method starts from a review literature about magic and antimagic labelings, in particular *(super) (a, d)-H-antimagic total labeling*, then continues by constructing the labeling on corona product of two paths. This construction produces several conjectures whose validity will be proven, where the proof process will utilize several techniques in k -balanced multisets and (k, δ) -anti balanced multisets. The given conjecture will be stated as a corollary, lemma, or theorem, if it is proven to be true. Otherwise, the research will be repeated by reconsidering the conjecture, the construction of the labeling, or the proof method used (Prihandini & Adawiyah, 2022).

C. RESULT AND DISCUSSION

Let P_n and P_m be two paths of order $n \geq 3$ and $m \geq 3$, respectively. In this paper, we consider corona product $P_n \odot P_m$. Let

$$V(P_n \odot P_m) = \{v_i \mid i = 1, 2, \dots, n\} \cup \{v_i^j \mid i = 1, 2, \dots, n, j = 1, 2, \dots, m\},$$

and

$$E(P_n \odot P_m) = \{v_i v_{i+1} \mid i = 1, 2, \dots, n-1\} \cup \{v_i v_i^j \mid i = 1, 2, \dots, n, j = 1, 2, \dots, m-1\} \cup \{v_i^j v_i^{j+1} \mid i = 1, 2, \dots, n, j = 1, 2, \dots, m-1\}.$$

Hence, we have $|V(P_n \odot P_m)| = mn + n$ and $|E(P_n \odot P_m)| = 2mn - 1$. Figure 1 below illustrates the general form of corona product $P_n \odot P_m$.




Figure 1. Corona product $P_n \odot P_m$

In this section, we define $\sum X = \sum_{x \in X} x$ and $\{a\} \cup \{a, b\} = \{a, a, b\}$. This section is divided into three subsections. In Subsection 3.1 and 3.2., we provide the definition of k -balanced multisets and k -anti balanced multisets, respectively, and also some useful lemmas that will be used to prove our main results about the super (a, d) - $P_2 \odot P_m$ -antimagic total labeling of $P_n \odot P_m$ in Subsection 3.

1. k -balanced multisets

A *multiset* is a set that allows the same elements to appear more than once within the set. One technique used to partition a multiset was initially introduced by (Maryati et al., 2013). Let $k \in \mathbb{N}$ and Y be a multiset containing positive integers. We said the multiset Y as k -balanced if there exists k subsets of Y , namely $Y_1, Y_2, \dots, Y_k$, so that for each $i = 1, 2, \dots, k$, the subset satisfies $|Y_i| = \frac{|Y|}{k}$, $\sum Y_i = \frac{\sum Y}{k} \in \mathbb{N}$, and $\cup_{i=1}^k Y_i = Y$. Such subset Y_i for each $i = 1, 2, \dots, k$ is called a *balanced subset* of Y . The following lemma is one of the k -balanced multisets.

Lemma 1. (Maryati et al., 2013) Let $x, y, k \in \mathbb{Z}$ so that $1 \leq x \leq y$ and $k > 1$. If $X = [x, y]$ and $|X|$ is a multiple of $2k$, then X is k -balanced.

Proof. For each $i = 1, 2, \dots, k$, define $X_i = \{\alpha_j^i \mid 1 \leq j \leq \frac{|X|}{k}\}$ where

$$\alpha_j^i = \begin{cases} x - 1 + (j - 1)k + i, & \text{for odd } j; \\ x + jk - i, & \text{for even } j. \end{cases}$$

It is not hard to check that for each $i = 1, 2, \dots, k$, we have $|X_i| = \frac{|X|}{k}$, $\cup_{i=1}^k X_i = X$, and $\sum X_i = \frac{|X|}{2k}(x + y)$. Therefore, X is k -balanced. ■

The following two corollaries are an immediate consequence of Lemma 1, which will be used to prove our main results.

Corollary 1. Let $m, n \in \mathbb{Z}$ with $m, n \geq 3$. Let $x = n + 1, y = 2mn + n$, and $k = mn$. If $B = [n + 1, 2mn + n]$ and $|B|$ is a multiple of $2mn$, then B is mn -balanced.

Proof. Note that $|B| = 2mn$. For each $i = 1, 2, \dots, mn$, define $B_i = \{b_j^i | 1 \leq j \leq \frac{2mn}{mn}\}$ where

$$b_j^i = \begin{cases} n + i, & \text{for } j = 1; \\ n + 2mn + 1 - i, & \text{for } j = 2. \end{cases}$$

it is not hard to check that for each $i = 1, 2, \dots, mn$, we have $|B_i| = 2$, $\cup_{i=1}^{mn} B_i = B$, and $\sum B_i = 2mn + 2n + 1$. Therefore, B is mn -balanced. ■

Corollary 2. Let $m, n \in \mathbb{Z}$ with $m, n \geq 3$ and m is odd. Let $x = 2mn + 2n, y = 3mn + n - 1$, and $k = n$. If $C = [2mn + 2n, 3mn + n - 1]$ and $|C|$ is a multiple of $2n$, then C is n -balanced.

Proof. Note that $|C| = (m - 1)n$. For each $i = 1, 2, \dots, n$, define $C_i = \{c_j^i | 1 \leq j \leq m - 1\}$ where

$$c_j^i = \begin{cases} 2mn + 2n - 1 + (j - 1)n + i, & \text{for odd } j; \\ 2mn + 2n + jn - i, & \text{for even } j. \end{cases}$$

it is not hard to check that for each $i = 1, 2, \dots, n$, we have $|C_i| = m - 1$, $\cup_{i=1}^n C_i = C$, and $\sum C_i = \frac{5m^2n - 2mn - m - 3n + 1}{2}$. Therefore, C is n -balanced. ■

2. (k, δ) -anti balanced multisets

Another technique of partitioning a multiset was introduced by Maryati et al. in (Maryati et al., 2013). Let $k \in \mathbb{N}$ and Y be a multiset containing positive integers. The multiset Y is said to be (k, δ) -anti balanced if there are k subsets of Y , namely $Y_1, Y_2, \dots, Y_k$, so that for each $i = 1, 2, \dots, k$, $|Y_i| = \frac{|Y|}{k}$, $\cup_{i=1}^k Y_i = Y$, and for each $i = 1, 2, \dots, k - 1$, $\sum Y_{i+1} - \sum Y_i = \delta$.

We now provide two lemmas about (k, δ) -anti balanced multisets which will be used to prove our main results.

Lemma 2. Let $m, n \in \mathbb{Z}$ with $m, n \geq 3$. Then $K = [1, n] \cup [2, n - 1] \cup [2mn + n + 1, 2mn + 2n - 1]$ is $(n - 1, 1)$ -anti balanced.

Proof. Note that $|K| = 3(n - 1)$. For each $i = 1, 2, \dots, n - 1$, define $K_i = \{a_i, b_i, c_i\}$ where

$$\begin{aligned} a_i &= i; \\ b_i &= i + 1; \\ c_i &= 2mn + 2n - i. \end{aligned}$$

It is not hard to check for each $i = 1, 2, \dots, n - 1$, $|K_i| = 3$, $\cup_{i=1}^{n-1} K_i = K$, $\sum K_i = 2mn + 2n + 1 + i$, and for each $i = 1, 2, \dots, n - 2$, $\sum K_{i+1} - \sum K_i = 1$. Therefore, K is $(n - 1, 1)$ -anti balanced. ■

Lemma 3. Let $m, n \in \mathbb{Z}$ with $m, n \geq 3$ and m is even. Then

$$\begin{aligned} L &= [2mn + 2n, 3mn + n - 1] \cup \left\{ 2mn + 2n + 2n(j - 1) + i \mid i = 1, 2, \dots, n - 2, j = 1, 2, \dots, \frac{m}{2} \right\} \\ &\quad \cup \left\{ 2mn + 4n + 2n(j - 1) - i - 1 \mid i = 1, 2, \dots, n - 2, j = 1, 2, \dots, \frac{m}{2} - 1 \right\} \end{aligned}$$

is $(n - 1, 2)$ -anti balanced.

Proof. Note that $|L| = 2(m-1)(n-1)$. For each $i = 1, 2, \dots, n-1$, we first define $L_i = \{a_j^i, b_j^i, c_j^i, d_j^i\}$ where

$$a_j^i = 2mn + 2n + i - 1 + n(j-1), \text{ for odd } j = 1, 2, \dots, m-1;$$

$$b_j^i = 2mn + 2n + i + n(j-1), \text{ for odd } j = 1, 2, \dots, m-1;$$

$$c_j^i = 2mn + 2n + nj - i, \text{ for even } j = 1, 2, \dots, m-1;$$

$$d_j^i = 2mn + 2n + nj - i - 1, \text{ for even } j = 1, 2, \dots, m-1.$$

It is not hard to check for each $i = 1, 2, \dots, n-1$, $|L_i| = 2(m-1)$, $\cup_{i=1}^{n-1} L_i = L$, $\sum L_i = 5m^2n - 2mn - m - 4n + 2i + 1$, and for each $i = 1, 2, \dots, n-2$, $\sum L_{i+1} - \sum L_i = 2$. Therefore, L is $(n-1, 2)$ -anti balanced. ■

3. Super (a, d) - $P_2 \odot P_m$ -antimagic total labeling of $P_n \odot P_m$

We now ready to provide our main results. In this subsection, we show that $P_n \odot P_m$ for $m, n \geq 3$ is super (a, d) - $P_2 \odot P_m$ -antimagic where $d = 1$ if m is odd and $d = 3$ if m is even.

Theorem 1. Let $m, n \in \mathbb{Z}$ with $m, n \geq 3$ and m is odd. Then the corona product $P_n \odot P_m$ is super $(9m^2n + 4mn + m - n + 3, 1)$ - $P_2 \odot P_m$ -antimagic.

Proof. Let $\lambda = [1, 3mn + n - 1]$ denotes set of labels for all vertices and edges of $P_n \odot P_m$. Partition λ into three subsets, $\lambda = A \cup B \cup C$, where $A = [1, n] \cup [2mn + n + 1, 2mn + 2n - 1]$, $B = [n + 1, 2mn + n]$ and $C = [2mn + 2n, 3mn + n - 1]$. We first define a function $f: V(P_n \odot P_m) \cup E(P_n \odot P_m) \rightarrow [1, 3mn + n - 1]$ as follows.

- (i) Define $K = A \cup [2, n-1]$. According to Lemma 2, set K is $(n-1, 1)$ -anti balanced where for each $i = 1, 2, \dots, n-1$, $K_i = \{a_i, b_i, c_i\}$ and $\sum K_i = 2mn + 2n + 1 + i$. We label vertices v_i and edges $v_i v_{i+1}$ by using the elements of K as follows.

$$f(v_i) = \begin{cases} \min K_i, & \text{for } i = 1, 2, \dots, n-1; \\ n, & \text{for } i = n; \end{cases}$$

$$f(v_i v_{i+1}) = \max K_i, \text{ for } i = 1, 2, \dots, n-1.$$

- (ii) According to Corollary 1, set B is mn -balanced where for each $i = 1, 2, \dots, mn$, $B_i = \{b_j^i | 1 \leq j \leq 2\}$ and $\sum B_i = 2mn + 2n + 1$. We label vertices v_i^j and edges $v_i^j v_i^{j+1}$ by using the elements of B as follows.

$$f(v_i^j) = \min B_{j+m(i-1)}, \text{ for } i = 1, 2, \dots, n, j = 1, 2, \dots, m;$$

$$f(v_i^j v_i^{j+1}) = \max B_{j+m(i-1)}, \text{ for } i = 1, 2, \dots, n, j = 1, 2, \dots, m.$$

- (iii) According to Corollary 2, set C is n -balanced where for each $i = 1, 2, \dots, n$, $C_i = \{c_j^i | 1 \leq j \leq m-1\}$ and $\sum C_i = \frac{5m^2n - 2mn - m - 3n + 1}{2}$. We label edges $v_i^j v_i^{j+1}$ by using the elements of C as follows.

$$f(v_i^j v_i^{j+1}) = c_j^i, \text{ for } i = 1, 2, \dots, n, j = 1, 2, \dots, m-1.$$

It is easy to see that the function f defined in (i)-(iii) is a bijective function from $V(P_n \odot P_m) \cup E(P_n \odot P_m)$ to $\{1, 2, \dots, 3mn + n - 1\}$ where $f(V(P_n \odot P_m)) = \{1, 2, \dots, mn + n\}$.

Now, observe that $P_n \odot P_m$ contains $n - 1$ subgraphs that isomorphic to $P_2 \odot P_m$. Then for each $i = 1, 2, \dots, n - 1$, the $P_2 \odot P_m$ -weight of the i -th subgraph $P_2 \odot P_m$ is

$$\begin{aligned} \omega(P_2 \odot P_m)_i &= \sum K_i + 2m \sum B_i + 2 \sum C_i \\ &= (2mn + 2n + 1 + i) + 2m(2mn + 2n + 1) + 2 \left(\frac{5m^2n - 2mn - m - 3n + 1}{2} \right) \\ &= 9m^2n + 4mn + m - n + 2 + i \end{aligned}$$

Since $\omega(P_2 \odot P_m)_2 = 9m^2n + 4mn + m - n + 3$ and $\omega(P_2 \odot P_m)_{i+1} - \omega(P_2 \odot P_m)_i = 1$ for each $i = 1, 2, \dots, n - 2$, we obtain that f is a super $(9m^2n + 4mn + m - n + 3, 1)$ - $P_2 \odot P_m$ -antimagic total labeling of $P_n \odot P_m$. Thus, $P_n \odot P_m$ is super $(9m^2n + 4mn + m - n + 3, 1)$ - $P_2 \odot P_m$ -antimagic. ■

Figure 2 below illustrates a super $(282, 1)$ - $P_2 \odot P_3$ -antimagic total labeling of $P_3 \odot P_3$.




Figure 2. A super $(282, 1)$ - $P_2 \odot P_3$ -antimagic total labeling of $P_3 \odot P_3$.

Theorem 2. Let $m, n \in \mathbb{Z}$ with $m, n \geq 3$ and m is even. Then the corona product $P_n \odot P_m$ is super $(9m^2n + 4mn + m - 2n + 5, 3)$ - $P_2 \odot P_m$ -antimagic.

Proof. Let $\lambda = [1, 3mn + n - 1]$ denotes a set of labels for all vertices and edges of $P_n \odot P_m$. In this proof, we also partition λ into three subsets, $\lambda = A \cup B \cup C$, where $A = [1, n] \cup [2mn + n + 1, 2mn + 2n - 1]$, $B = [n + 1, 2mn + n]$ and $C = [2mn + 2n, 3mn + n - 1]$. We first define a function $f: V(P_n \odot P_m) \cup E(P_n \odot P_m) \rightarrow [1, 3mn + n - 1]$ as follows.

- (i) Define $K = A \cup [2, n - 1]$. According to Lemma 2, set K is $(n - 1, 1)$ -anti balanced where for each $i = 1, 2, \dots, n - 1$, $K_i = \{a_i, b_i, c_i\}$ and $\sum K_i = 2mn + 2n + 1 + i$. We label vertices v_i and edges $v_i v_{i+1}$ by using the elements of K as follows.

$$f(v_i) = \begin{cases} \min K_i, & \text{for } i = 1, 2, \dots, n - 1; \\ n, & \text{for } i = n; \end{cases}$$

$$f(v_i v_{i+1}) = \max K_i, \text{ for } i = 1, 2, \dots, n - 1.$$

- (ii) According to Corollary 1, set B is mn -balanced where for each $i = 1, 2, \dots, mn$, $B_i = \{b_j^i | 1 \leq j \leq 2\}$ and $\sum B_i = 2mn + 2n + 1$. We label vertices v_i^j and edges $v_i^j v_i^l$ by using the elements of B as follows.

$$f(v_i^j) = \min B_{j+m(i-1)} \text{ , for } i = 1, 2, \dots, n, j = 1, 2, \dots, m;$$

$$f(v_i v_i^j) = \max B_{j+m(i-1)} \text{ , for } i = 1, 2, \dots, n, j = 1, 2, \dots, m.$$

(iii) Define

$$L = [2mn + 2n, 3mn + n - 1] \cup \left\{ 2mn + 2n + 2n(j-1) + i \mid i = 1, 2, \dots, n-2, j = 1, 2, \dots, \frac{m}{2} \right\} \cup \left\{ 2mn + 4n + 2n(j-1) - i - 1 \mid i = 1, 2, \dots, n-2, j = 1, 2, \dots, \frac{m}{2} - 1 \right\}.$$

(iv) According to Lemma 3, set L is n is $(n-1, 2)$ -anti balanced where for each $i = 1, 2, \dots, n-1$, $L_i = \{a_i^j, b_i^j, c_i^j, d_i^j\}$ and $\sum L_i = 5m^2n - 2mn - m - 4n + 2i + 1$. We label edges $v_i^j v_i^{j+1}$ by using the elements of L as follows.

$$f(v_i^j v_i^{j+1}) = \begin{cases} 2mn + 2n + i - 1 + n(j-1), & \text{for odd } j = 1, 2, \dots, m-1; \\ 2mn + 2n + nj - i, & \text{for even } j = 1, 2, \dots, m-1. \end{cases}$$

It is easy to see that the function f defined in (i)-(iii) is a bijective function from $V(P_n \odot P_m) \cup E(P_n \odot P_m)$ to $\{1, 2, \dots, 3mn + n - 1\}$ where $f(V(P_n \odot P_m)) = \{1, 2, \dots, mn + n\}$. Now, observe that $P_n \odot P_m$ contains $n-1$ subgraphs that isomorphic to $P_2 \odot P_m$. Then for each $i = 1, 2, \dots, n-1$, the $P_2 \odot P_m$ -weight of the i -th subgraph $P_2 \odot P_m$ is

$$\begin{aligned} \omega(P_2 \odot P_m)_i &= \sum K_i + 2m \sum B_i + \sum L_i \\ &= (2mn + 2n + 1 + i) + 2m(2mn + 2n + 1) + 5m^2n - 2mn - m - 4n + 2i + 1 \\ &= 9m^2n + 4mn + m - 2n + 3i + 2 \end{aligned}$$

Since $\omega(P_2 \odot P_m)_1 = 9m^2n + 4mn + m - 2n + 5$ and $\omega(P_2 \odot P_m)_{i+1} - \omega(P_2 \odot P_m)_i = 3$ for each $i = 1, 2, \dots, n-2$, we obtain that f is a super $(9m^2n + 4mn + m - 2n + 5, 3) \cdot P_2 \odot P_m$ -antimagic total labeling of $P_n \odot P_m$. Thus, $P_n \odot P_m$ is super $(9m^2n + 4mn + m - 2n + 5, 3) \cdot P_2 \odot P_m$ -antimagic. ■

Figure 3 below illustrates a super $(483, 3) \cdot P_2 \odot P_4$ -antimagic total labeling of $P_3 \odot P_4$.




Figure 3. A super $(483, 3) \cdot P_2 \odot P_4$ -antimagic total labeling of $P_3 \odot P_4$.

D. CONCLUSION AND SUGGESTIONS

We have shown that the corona product $P_n \odot P_m$ for $m, n \geq 3$ is super $(9m^2n + 4mn + m - n + 3, 1) - P_2 \odot P_m$ -antimagic if m is odd and super $(9m^2n + 4mn + m - 2n + 5, 3) - P_2 \odot P_m$ -antimagic if m is even. The values of d obtained in paper are limited, hence it is interesting for the readers to studying the super $(a, d) - P_2 \odot P_m$ -antimagic total labeling of $P_n \odot P_m$ for the remaining possible values of d . Further, the readers can also consider to studying the super $(a, d) - P_r \odot P_m$ -antimagic total labeling of $P_n \odot P_m$ for other possible values of r .

ACKNOWLEDGEMENT

The research for this paper was supported financially by PNB Program from Universitas Mataram.

REFERENCES

- Agustin, I. H., Prihandini, R. M., & Dafik. (2019). $P_2 \triangleright H$ -super antimagic total labeling of comb product of graphs. *AKCE International Journal of Graphs and Combinatorics*, 16(2), 163–171. <https://doi.org/10.1016/j.akcej.2018.01.008>
- Anjancylu, G. S. G. N., Vijayabarathi, A., & Liu, L. (2015). Super edge-magic sequence of maximal outer planer graph and its characteristics. *Cogent Mathematics*, 2(1), 1123340. <https://doi.org/10.1080/23311835.2015.1123340>
- Bača, M., Miller, M., Ryan, J., & Semaničová-Feňovčíková, A. (2019). *Magic and Antimagic Graphs* (1st ed., Vol. 60). Springer International Publishing. <https://doi.org/10.1007/978-3-030-24582-5>
- Chang, F., Chen, H., & et al. (2019). Shifted-Antimagic Labelings for Graphs. *Graphs and Combinatorics*, 37(2021), 1065–1082.
- Diestel, R. (2017). *Graph Theory* (5th ed., Vol. 173). Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-662-53622-3>
- Enomoto, H., Llado, A. S., Nakamigawa, T., & Ringel, G. (1998). SUPER EDGE-MAGIC GRAPHS. *SUT Journal of Mathematics*, 34(2). <https://doi.org/10.55937/sut/991985322>
- Gallian, J. A. (2022). A Dynamic Survey of Graph Labeling. In the electronic journal of combinatorics.
- Gutiérrez, A., & Lladó, A. (2005). Magic coverings. *Journal of Combinatorial Mathematics and Combinatorial Computing*, 55, 43.
- Hasni, R., Tarawneh, I., & Husin, M. Na. (2022). A Remark On The Edge Irregularity Strength Of Corona Product Of Two Paths. *Journal of Mathematical Sciences and Informatics*, 2(1), 51–58. <https://doi.org/10.46754/jmsi.2022.06.005>
- Hendy, H., Sugeng, K. A., & Salman, A. N. M. (2018). An H -super magic decompositions of the lexicographic product of graphs. *AIP Conference Proceedings*, 2023. <https://doi.org/10.1063/1.5064190>
- Hendy, H., Sugeng, K. A., Salman, A. N. M., & Ayunda, N. (2018). Another H-super magic decompositions of the lexicographic product of graphs. *Indonesian Journal of Combinatorics*, 2(2), 72. <https://doi.org/10.19184/ijc.2018.2.2.2>
- Hussain, M., Baskoro, E. T., Ali, K., & Hussain, M. (2012). On super antimagic total labeling of Harary graph. <https://www.researchgate.net/publication/228458682>
- Inayah, N., Salman, A. N. M., & Simanjuntak, R. (2009). On (a, d) -H-antimagic coverings of graphs.
- Inayah, N., Simanjuntak, R., Salman, A. N. M., & Syuhada, K. I. A. (2013). Super (a, d) -H-antimagic total labelings for shackles of a connected graph H. In *Australasian Journal Of Combinatorics* (Vol. 57).

- Indunil, minusha, & Perera.A.A.I. (2022). Graceful Labeling of Chain Graphs with Pendants. *International Conference On Business Innovations 2022*, 595–598.
- Kotzig, A., & Rosa, A. (1970). Magic Valuations of Finite Graphs. *Canadian Mathematical Bulletin*, 13(4), 451–461. <https://doi.org/10.4153/cmb-1970-084-1>
- Liu, J. B., Afzal, H. U., & Javaid, M. (2020). Computing Edge Weights of Magic Labeling on Rooted Products of Graphs. *Mathematical Problems in Engineering*, 2020. <https://doi.org/10.1155/2020/2160104>
- Llado A, & Moragas J. (2007). Cycle-Magic Graphs. *DiscreteMath*, 307, 2925–2933.
- Martin Bača, Martin Bača, Rinovia Simanjuntak, & Rinovia Simanjuntak. (2006). Super edge-antimagic labelings of the generalized Petersen graph $P(n, (n-1)/2)$. *Utilitas Mathematica*, 70, 119–127.
- Maryati, T. K., Salman, A. N. M., & Baskoro, E. T. (2013). Supermagic coverings of the disjoint union of graphs and amalgamations. *Discrete Mathematics*, 313(4), 397–405. <https://doi.org/10.1016/j.disc.2012.11.005>
- Ngurah, A. A. G., Salman, A. N. M., & Susilowati, L. (2010). H-supermagic labelings of graphs. *Discrete Mathematics*, 310(8), 1293–1300. <https://doi.org/10.1016/j.disc.2009.12.011>
- Nurvazly, D. E., Chasanah, S. L., & Wiranto, A. R. (2022). Variations of graceful labelling of subgraph of millipede graph. *AIP Conference Proceedings*, 050018. <https://doi.org/10.1063/5.0105447>
- Palanivelu, C., & Neela, N. (2019). Super (a, d)-Edge Antimagic Total Labeling of Union of Stars. In *International Journal of Applied Engineering Research* (Vol. 14, Issue 9). <http://www.ripublication.com>
- Permata Sari, A., Sri Martini, T., & Yugi Kurniawan, V. (2019). Super (a, d)-H-antimagic total labeling of edge corona product on cycle with path graph and cycle with cycle graph. *Journal of Physics: Conference Series*, 1306(1). <https://doi.org/10.1088/1742-6596/1306/1/012006>
- Prihandini, R. M., & Adawiyah, R. (2022). On Super (a,d)-Edge Antimagic Total Labeling of Some Generalized Shackle of Fan Graph. In *International Journal of Academic and Applied Research* (Vol. 6). www.ijeais.org/ijaar
- Prihandini, R. M., & Adawiyah, R. (2023). On Super $(3n+5,2)$ - Edge Antimagic Total Labeling And It's Application To Construct Hill Chiper Algorithm. *BAREKENG: Jurnal Ilmu Matematika Dan Terapan*, 17(1), 0029–0036. <https://doi.org/10.30598/barekengvol17iss1pp0029-0036>
- Raheem, A., Baig, A. Q., & Javaid, M. (2014). On Super (a, d)-Edge-Antimagic Total Labeling of a Class of Trees. In *International J.Math. Combin* (Vol. 4).
- Rajeswari, R., Aruna Sakthi, K., & Meenakumari, N. (2021). Product-Set Labeling of Some Zero-Divisor Graphs and its Line Graph. *Journal of Physics: Conference Series*, 1947(1). <https://doi.org/10.1088/1742-6596/1947/1/012025>
- Sandariria, H., Roswitha, M., & Kusmayadi, T. A. (2017). H-Supermagic Labeling on Coronation of Some Classes of Graphs with a Path. *Journal of Physics: Conference Series*, 855(1). <https://doi.org/10.1088/1742-6596/855/1/012042>
- Simanihuruk, M., Kusmayadi, T. A., Swita, B., Romala, M., & Damanik, F. (2021). A Conjecture on Super Edge-Magic Total Labeling of 4-Cycle Books. *International Journal of Mathematics and Mathematical Sciences*, 2021. <https://doi.org/10.1155/2021/8483926>
- Simanjuntak, R., Bertault, F., & Miller, M. (2000). Two new (a, d)-antimagic graph labelings. *Proc. of Eleventh Australasian Workshop on Combinatorial Algorithms*, 11, 179–189.
- Lakshmi, S., & Sagayakavitha, D. (2018). Edge-magic total labeling of some graphs. *International Journal of Mathematics Trends and Technology (IJMTT)*, 1, 101–103.
- Taimur, A., Numan, M., Ali, G., Mumtaz, A., & Semaničová-Feňovčíková, A. (2018). Super (a, d)-H-antimagic labeling of subdivided graphs. *Open Mathematics*, 16(1), 688–697. <https://doi.org/10.1515/math-2018-0062>

- Ulfatihah, R., Roswitha, M., & Kusmayadi, T. A. (2017). H-Supermagic Labeling on Shrubs Graph and $L_m \odot P_n$. Journal of Physics: Conference Series, 855(1). <https://doi.org/10.1088/1742-6596/855/1/012055>
- Zeen El Deen, M. R. (2019). Edge even graceful labeling of some graphs. Journal of the Egyptian Mathematical Society, 27(1). <https://doi.org/10.1186/s42787-019-0025-x>

Artikel Bela

ORIGINALITY REPORT

9%

SIMILARITY INDEX

7%

INTERNET SOURCES

9%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1

www.jocm.us

Internet Source

1%

2

dokumen.pub

Internet Source

1%

3

Submitted to UIN Syarif Hidayatullah Jakarta

Student Paper

1%

4

R. Rajeswari, K. Aruna Sakthi, N. Meenakumari. "Product-Set Labeling of Some Zero-Divisor Graphs and its Line Graph", Journal of Physics: Conference Series, 2021

Publication

1%

5

Submitted to Universitas Negeri Surabaya
The State University of Surabaya

Student Paper

1%

6

ejournal.unesa.ac.id

Internet Source

1%

7

digilib.unila.ac.id

Internet Source

1%

8

eprints.unipdu.ac.id

Internet Source

1%

9

researchonline.federation.edu.au

Internet Source

1 %

10

G.S.G.N. Anjaneyulu, A. Vijayabarathi, Lishan Liu. "Super edge-magic sequence of maximal outer planer graph and its characteristics", Cogent Mathematics, 2015

Publication

Exclude quotes Off

Exclude matches < 1%

Exclude bibliography On